

RELATIVE GROWTH OF *GONIOPSIS CRUENTATA* (CRUSTACEA, BRACHYURA, GRAPSIDAE), ON THE UBATUBA REGION, SÃO PAULO, BRAZIL

Valter J. Cobo¹
Adilson Fransozo²

ABSTRACT

Relative growth analyses had specially focused the determination of size at sexual maturity by means of evidencing the puberty molt. Description of relative growth patterns in *Goniopsis cruentata* (Latreille, 1803), and the estimation of the onset of morphological sexual maturity was studied. Monthly surveys were conducted from June 1992 to July 1993. Crabs were hand captured by three people during 1 h sample effort, in an estuarine area in the Ubatuba region, São Paulo, Brazil (23°29'24"S; 45°10'12"W). Measurements of carapace width and length (CW and CL), abdomen width (AW), chelar propodus height (CPH) and gonopod length (GL) in males, were recorded. Regressions were performed according to the general relative growth formula $y = a \cdot x^b$. A student's t-test was applied to evaluate statistically the allometric growth constant (b). Different population groups were considered: young and adult males, young, pre-pubertal and adult females. The morphometric relations of the carapace showed a tendency to isometry. In females the abdominal width grow in positive allometry: young $b = 1.35$; pre-pubertal $b = 1.58$ and adult $b = 1.38$. In males, the growth of gonopod showed a positive allometry, for young ($b = 1.64$) and an isometric relation in adult phase ($b = 0.957$). This trend reflects an energetic budget allocation biased towards reproduction rather than growth purposes after the puberty molt.

KEYWORDS. Grapsidae, *Goniopsis*, allometric growth, puberty molt.

INTRODUCTION

Literature devoted to relative growth in Brachyura approach the mathematical theory and the principles of allometry (HUXLEY, 1950; TEISSIER, 1960; HARTNOLL, 1982), besides the aspects related to the determination of size at the onset of sexual maturity (WEYMOUTH & MACKAY, 1934; GORE & SCOTTO, 1983; HAEFNER, 1990). Observed variability in growth patterns between mature and immature growth phases and within sexes are a result of relative development

1. Departamento de Biologia, Universidade de Taubaté (UNITAU), CEP 12.020-270 Taubaté, São Paulo, Brasil.

2. Departamento de Zoologia, Instituto de Biociências e Centro de Aqüicultura, UNESP, CEP 18618-000 Botucatu, São Paulo, Brasil.

differences of certain structures (HARTNOLL, 1974).


Relative growth studies of brachyurans from São Paulo State Coast, emphasized the attainment of sexual maturity, e.g. PINHEIRO & FRANSOZO (1993) who studied the portunid *Arenaeus cribrarius* (Lamark, 1818), MANTELLATO & FRANSOZO (1994) on *Hepatus pudibundus* (Herbst, 1785), HIYODO & FRANSOZO (1995) regarding the spider crab *Acantonyx scutiformis* (Dana, 1851) and SANTOS *et al.* (1995) on *Portunus spinimanus* Latreille, 1819.

The present study intended to describe the relative growth patterns of *Goniopsis cruentata* (Latreille, 1803). Also, the transition interval between immature and mature growth phases in allometric relationships were assessed, as an attempt to estimate the size at the onset of morphological maturity.

MATERIAL AND METHODS

According to MELO (1996) the geographical range of *Goniopsis cruentata* is restricted to the Occidental Atlantic, from Florida (USA), Bermuda, Gulf of Mexico, Central America, South America; in Brazil, the species was recorded from Fernando de Noronha to Santa Catarina State.

During this study, monthly samples from June 1992 to July 1993 were conducted in a mangrove area, located between Comprido and Escuro rivers ($23^{\circ} 29' 24''$ S, $45^{\circ} 10' 12''$ W), Fortaleza Bay, Ubatuba, SP. Crabs were hand captured by three people during 1 h sample effort collections, and placed in individual bags which were frozen afterwards and defrosted prior to laboratory analyses. A vernier caliper was used to measure the carapace width (CW), length (CL), abdomen width (AW) and chelar propodus height (CPH) in all crabs. The gonopod length of males was measured with the aid of a stereomicroscope provided with a camera lucida (figs. 1-5).




Figs. 1-5 . *Goniopsis cruentata*, measurements: 1, length and width (L,W) of carapace; 2, width of female abdome; 3, width of male abdome; 4, height and width (H,W) of propodus; 5, lenght of gonopod.

The general relative growth formula, $y = a \cdot x^b$ (HUXLEY, 1950; TEISSIER, 1960; HARTNOLL, 1974, 1982) was used, in which (y) represents the dimension under analysis (dependent variable), (a) is the intersection, (b) the allometric growth constant and (x) a dimension regarded as size (independent variable). The relative growth coefficient, (b) was statistically evaluated in all resulting relationships by means of a student's t-test at the 5% significance level.

RESULTS

A total of 474 crabs were obtained during the study period. In order to analyze the relative growth relationships, males were differentiated in young and adult, and females were distinguished in young, pre-pubertal and adult, regarding their abdominal external morphology (figs. 6-10, tab. I).

Carapace growth, as expressed in the CW x CL relationship, is isometric and continuous in all relationships performed, suggesting that a single allometric relationship must be applied for each sex. AW x CW relationships evidenced a positive allometric growth in all female groups (fig.11). In males, a positive allometry was only recorded in the young, while adults presented an isometric abdominal growth.



Figs. 6-10. *Goniopsis cruentata*, shape of abdoles: 6, young male; 7, pre-pubertal female; 8, young female; 9, adult male; 10, adult female.

Table I. Regression analysis of the morphometric data of *Goniopsis cruentata*, on the Ubatuba region, SP/ Brazil. (AM= adult male; YM= young male; TM= total male; AF= adult female; PPF= pre-pubertal female; YF= young female and TF= total female) (AW= abdome width; PH= propodus height; GL gonopod length; CL= carapace length. (=) isometry (+) positive allometry.

Variable	Group	N	$y=a.x^b$	$\ln y=\ln a+b.\ln x$	r^2	Allometry
AW	AM	206	AW=0.442 CW ^{0.964}	$\ln AW=-0.816+0.964 \ln CW$	0.96	=
	YM	23	AW= 0.182 CW ^{1.50}	$\ln AW= -1.70+1.25 \ln CW$	0.92	+
	TM	229	AW= 0.321 CW ^{1.06}	$\ln AW= -1.14+1.06 \ln CW$	0.97	=
	AF	207	AW= 0.165 CW ^{1.38}	$\ln AW= -1.80+1.38 \ln CW$	0.83	+
	YF	13	AW= 0.147 CW ^{1.35}	$\ln AW= -1.92+1.35 \ln CW$	0.91	+
	PPF	25	AW= 0.078 CW ^{1.58}	$\ln AW= -2.54+1.58 \ln CW$	0.83	+
	TF	245	AW= 0.106 CW ^{1.50}	$\ln AW= -2.25+1.50 \ln CW$	0.93	+
PH	AM	206	PH= 0.058 CW ^{1.51}	$\ln PH= -2.85+1.51 \ln CW$	0.90	+
	YM	23	PH= 0.138 CW ^{1.21}	$\ln PH= -1.98+1.21 \ln CW$	0.95	+
	TM	229	PH= 0.365 CW ^{1.46}	$\ln PH= -2.67+1.46 \ln CW$	0.94	+
	AF	232	PH= 0.134 CW ^{1.22}	$\ln PH= -2.00+1.22 \ln CW$	0.83	+
	YF	13	PH= 0.210 CW ^{1.03}	$\ln PH= -1.56+1.03 \ln CW$	0.93	=
	TF	245	PH= 0.127 CW ^{1.24}	$\ln PH= -2.06+1.24 \ln CW$	0.92	+
	GL	AM	206	GL=0.272 CW ^{0.957}	$\ln GL= -1.30+0.95 \ln CW$	0.94
YM		23	GL= 0.019 CW ^{1.64}	$\ln GL= -3.94+1.64 \ln CW$	0.76	+
TM		229	GL= 0.025 CW ^{1.63}	$\ln GL= -3.68+1.63 \ln CW$	0.82	+
CL	TM	229	CL= 0.648 CW ^{1.06}	$\ln CL= -0.434+1.06 \ln CW$	0.99	=
	TF	245	CL= 0.699 CW ^{1.05}	$\ln CL= -0.337+1.05 \ln CW$	0.99	=
	TOTAL	474	CL= 0.673 CW ^{1.06}	$\ln CL= -0.396+1.06 \ln CW$	0.99	=


A positive allometric growth was evidenced in the AP x LC relationship in both young and adult males. In females a positive allometry was suggested only in adults.

GL x AW was the relationship in which the male growth phase transition was most clearly identified, due to a marked positive allometry recorded in young crabs (fig. 12).

DISCUSSION

According to HARTNOLL (1974) and HAEFNER (1990), a certain general pattern of growth is ascribed for brachyurans. During the immature phase, males chelae experienced a high positive allometry and a noteworthy increase of their relative size at the puberty molt. Females chelae follow an isometric growth through all phases. In females, the abdomen undergone a high positive allometry in the immature phase and a remarkable increase at the puberty molt. In the case of males, an isometric growth is often observed.

The morphometric patterns obtained for the allometric relationships in *G. cruentata* are similar to those proposed by HARTNOLL (1974) for brachyuran crabs, except for some differences concerning the immature phase in males and the



Figs. 11-12. *Goniopsis cruentata*: 11, Relationship between abdomen width (AW) and carapace width (CW) in females (n=256) (Scatterplot of empirical data. (□) young females; (△) pre-pubertal females; (■) adult females). 12, Relationship between gonopod length (GL) and carapace width (CW) in males (n=268). Scatterplot of empirical data. (□) young males; (■) adult males.

mature phase in females. Chelipeds of *G. cruentata* adult females and abdomen of young males showed a positive allometric growth. Both trends differ from the patterns proposed by HARTNOLL (1974).

HARTNOLL's (1974) patterns of growth are largely based on sexual dimorphism in the Brachyura, mainly those related to chelipeds and abdomen. In the case of chelipeds, behavioral patterns are directly implied. Males show more intense territorial contentions, courtship and inter and intraspecific fights, as well demonstrated in genus *Uca* by CRANE (1957, 1958), *Aratus pisonii* (H. Milne Edwards, 1837) by WARNER (1970) and in *Goneplax rhomboides* (Linnaeus, 1758) by ABELLÓ & SARDÁ (1982).

The resemblance between chelar relative growth patterns in both sexes suggests that behavioral displays in males and females should be similar, which could explain the parity of the allometric values obtained. This supposition is supported by similar morphometric data obtained by other authors, e.g. *Arenaeus*

cribrarius, (PINHEIRO & FRANSOZO, 1993), *Hepatus pudibundus*, (MANTELATTO & FRANSOZO, 1994) and *Persephona lichtensteinii* Leach, 1817, *P. mediterranea* (Herbst, 1794) and *P. punctata* (Linnaeus, 1758) by HIYODO (1996).

Concerning the abdomen, its dimorphic growth in *Brachyura* is usually related to reproductive functions after the puberty molt. In this sense, young males of *G. cruentata* show a pattern of abdominal growth in which the concomitant transformation of gonopods is closely followed. By this means, the abdomen is continuously fitted to enclose and protect the gonopods, i.e. the appendages directly related to reproduction. In females, the main responsible factor influencing the allometric levels recorded is certainly the incubation of eggs.

The relationships $GL \times CW$ in males, and $AW \times CW$ in females, are surely those which more clearly evidence the size at the onset of sexual maturity, because they are directly related to reproduction, a decisive process for *G. cruentata* success. This sort of allometric relationships are commonly used for this purpose, e.g. HARTNOLL (1965) on *Aratus pisonii*; HALLEY (1969), who studied *Ocypode quadrata* (Fabricius, 1787); FINNEY & ABELE (1981) on *Trapezia ferruginea* Latreille, 1825; DAVIDSON & MARSDEN (1987) in the case of *Ovalipes catharus* (White, 1843); VANNINI & GHERARDI (1988) on the xanthids *Eriphia smithi* MacCleay, 1838.

The gonopod growth of *G. cruentata*, suggests an initial development of this structure which will undergo remarkable changes regarding shape and size at the puberty molt. This fact is reflected in the positive allometry observed during the immature phase and the maintenance of its proportion after the puberty molt, in order to efficiently accomplish the reproductive functions during the adult phase.

G. cruentata showed a proportional growth of carapace dimensions as verified by PINHEIRO (1993), who revised portunid allometric patterns. This relationship did not show any evidence which could suggest the size at maturation attainment, or indicated an eventual sexual dimorphism. From the analyses obtained, it can be concluded that the size at the onset of maturity in males and females is slightly different. It may occur between 20 and 24 mm CW in males and 18 and 29 mm CW in females.

Size ranges as those observed herein are common, and may be due to intrinsic plasticity of organisms or influenced by environmental factors such as temperature, pluviosity, photoperiod, food availability and others (CAMPBELL & EAGLES, 1983). Likewise, differences of size at sexual maturity between sexes are not unusual and have been verified in other species, e.g. *Trapezia ferruginea* studied by FINNEY & ABELE (1981); in *Sesarma reticulatum* (Say, 1817) by SEIPLE & SALMON (1987) and for *Callinectes ornatus* Ordway, 1863, by HAEFNER (1990). This feature is certainly associated to species-specific reproductive strategies, in which the available number of mature females is maximized.

It is admitted that the growth pattern of *G. cruentata* tends to a positive allometry during the immature phase of development, a trend that was not applied to all mature relationships. This fact suggests that differences in the allometric levels should be concentrated in the young-adult transition range. Therefore, energy partitioning should be biased towards reproduction after the puberty molt. In contrast, during the immature phase the energetic budget is mainly directed

towards the development of certain organs and/or structures, in order to attain an adequate functional size. After the puberty molt, energetic costs of reproduction increase, while growth through the successive instars will be restricted to the maintenance of acquired body proportions.

Acknowledgments. To the Conselho Nacional de Desenvolvimento Científico e Tecnológico - CNPq (proc.840.211/95-3). To Dr. Fernando Luiz Medina Mantelatto (Universidade de São Paulo) for revising the manuscript and providing valuable suggestions. To the NEBECC members for their help in field collections.

REFERENCES

- ABELLÓ, P. & SARDÁ, F. 1982. Nota sobre la morfometría del dimorfismo sexual en las pinzas de *Goneplax rhomboides* (Linnaeus, 1758) (Decapoda, Brachyura). **Investigación Pesq.**, Barcelona, **46** (2): 163 - 170.
- CAMPBELL, G.R. & EAGLES, M.D. 1983. Size at maturity and fecundity of rock crabs, *Cancer irroratus*, from the bay of Fundy and southwestern Nova Scotia. **Fish. Bull.**, Washington, **81**(2): 357 - 362.
- CRANE, J. 1957. Basic patterns of displays in fiddler crabs (Ocypodidae, Genus *Uca*). **Zoologica**, Stuttgart, **42**: 69 - 83.
- _____. 1958. Aspects of social behavior in fiddler crabs, with special reference to *Uca maracoani* (Latreille). **Zoologica**, Stuttgart, **43**: 113 - 130.
- DAVIDSON, R.J & MARSDEN, I.D. 1987. Size relationships and relative growth of the New Zealand swimming crab *Ovalipes catharus* (White, 1843). **J. Crust. Biol.**, San Antonio, **7** (2): 308 - 317.
- FINNEY, W.C. & ABELE, L.G. 1981. Allometric variation and sexual maturity in the obligate coral commensal *Trapezia ferrugina* Latreille (Decapoda, Xanthidae). **Crustaceana**, Leiden, **41** (2): 113 - 130.
- GORE, R.H. & SCOTTO, L.L. 1983. Studies on Decapod Crustacea from the Indian river region of Florida. XXV carapacial and abdominal allometry in five species of subtropical parthenopid crabs (Brachyura, Parthenopidae). **Crustaceana**, Leiden, **44** (1): 1 - 22.
- HAEFNER Jr., P. A. 1990. Morphometry and size at maturity of *Callinectes ornatus* (Brachyura, Portunidae) in Bermuda. **Bull. Mar. Sci.**, Coral Gables, **46** (2): 274 - 286.
- HALLEY, S. R. 1969. Relative growth and sexual maturity of the Texas ghost crab *Ocypode quadrata* (Fabr.) (Brachyura, Ocypodidae). **Crustaceana**, Leiden, **17**: 285 - 297.
- HARTNOLL, R. G. 1965. Notes on the marine grapsid crabs of Jamaica. **Proc. Linn. Soc. Lond.**, Lancaster, **176**: 113 - 147.
- _____. 1974. Variation in growth pattern between some secondary sexual characters in crabs (Decapoda, Brachyura). **Crustaceana**, Leiden, **27** (2): 131 - 136.
- _____. 1982. Growth. In: BLISS, D.E. ed. **The Biology of Crustacea, Embryology, Morphology, and Genetics**. New York, Academic. v.2, cap. 3, p. 11 - 196.
- HIYODO, C. M. 1996. **Biologia populacional do gênero *Persephona* Leach, 1817 (Crustacea, Decapoda, Leucosilidae) na região de Ubatuba, SP.** 131p. Tese (Doutorado em Ciências Biológicas), Zoologia, UNESP, Botucatu. [Não Publicada].
- HIYODO, C.M. & FRANSOZO, A. 1995. Relative growth of spider crab *Acanthonyx scutiformis* (Dana, 1851) (Crustacea, Decapoda, Majidae). **Arq. Biol. Tecnol.**, Curitiba, **38** (3): 969 - 981.
- HUXLEY, J.S. 1950. Relative growth and form transformation. **Proc. R. Soc. Lond.**, London, **137** (B): 465 - 469.
- MANTELATTO, F.L.M. & FRANSOZO, A. 1994. Crescimento relativo e dimorfismo sexual em *Hepatus pudibundus* (Herbst, 1785) (Decapoda, Brachyura) no litoral norte paulista. **Papéis Avuls. Zool.**, São Paulo, **39** (4): 33 - 48.
- MELO, G.A.S. 1996. **Manual de identificação dos Brachyura (caranguejos e siris) do litoral brasileiro.** São Paulo, Plêiade, FAPESP. 604 p.
- PINHEIRO, M.A.A. & FRANSOZO, A. 1993. Relative growth of the speckled swimming crab *Arenaeus cribrarius* (Lamarck, 1818) (Brachyura, Portunidae) near Ubatuba, State of São Paulo, Brazil. **Crustaceana**, Leiden, **65** (3): 377 - 389.
- SANTOS, S.; NEGREIROS-FRANZOZO M.L. & FRANSOZO, A. 1995. Morphometric relationships and maturation in *Portunus spinimanus* Latreille, 1819 (Crustacea, Brachyura, Portunidae). **Revta bras. Biol.**, Rio de Janeiro, **55**, (4): 545 - 553.
- SEIPLE, W. & SALMON, M. 1987. Reproductive, growth and life-history contrasts between two species of grapsid crabs, *Sesarma cinereum* and *Sesarma reticulatum*. **Mar. Biol.**, Berlin, **94**: 1 - 6

- TEISSIER, G., 1960. Relative growth. In: WATERMAN, T.H. ed. **The Physiology of crustacea. Metabolism and growth**. New York, Academic, v. 1, p. 537 - 560.
- VANNINI, M. & GHERARDI, F. 1988. Studies on the peblee crab *Eriphia smithi* MacCley, 1838 (Xanthoidea, Menippidae): Patterns of relative growth and population structure. **Trop. Zool.**, Florence, 1: 203 - 216.
- WARNER, G. F. 1970. Behavior of two species of grapsid crabs during intraespecific encounters. **Behavior**, Leiden, 1: 203 - 216.
- WEYMOUTH, F.W. & MACKAY, D.C.G. 1934. Relative growth in the edible crab *Cancer magister*. **Proc. Soc. exp. Biol. Med.**, New York, 31 (9): 1137 - 1139.

Recebido em 24.02.1997; aceito em 15.10.1997.